

WEST Search History

Hide Items

Restore


Clear

Cancel

DATE: Monday, January 31, 2005

Hide?	Set Name	Query	Hit Count
		<i>DB=EPAB; PLUR=YES; OP=OR</i>	
<input type="checkbox"/>	L4	WO-2004001004-A2.did.	1
		<i>DB=PGPB,USPT,USOC,EPAB,JPAB,DWPI; PLUR=YES; OP=OR</i>	
<input type="checkbox"/>	L3	L2 and tem	8
<input type="checkbox"/>	L2	croix.in.	192
<input type="checkbox"/>	L1	st-croix.in.	0

END OF SEARCH HISTORY



Entrez PubMed Nucleotide Protein Genome Structure OMIM PMC Journals Books

Search for

Limits Preview/Index History Clipboard Details

Show:

About Entrez

Text Version

☐ 1: Cancer Res. 2004 Dec 1;64(23):8507-11.

[Related Articles, Link](#)

Entrez PubMed

Overview

Help | FAQ

Tutorial

New/Noteworthy

E-Utilities

PubMed Services

Journals Database

MeSH Database

Single Citation Matcher

Batch Citation Matcher

Clinical Queries

LinkOut

Cubby

Related Resources

Order Documents

NLM Catalog

NLM Gateway

TOXNET

Consumer Health

Clinical Alerts

ClinicalTrials.gov

PubMed Central

Full text article at
concerres.aacrjournals.org

Identification of a binding partner for the endothelial cell surface proteins TEM7 and TEM7R.

Nanda A, Buckhaults P, Seaman S, Agrawal N, Boutin P, Shankara S, Nacht M, Teicher B, Stampfl J, Singh S, Vogelstein B, Kinzler KW, St Croix B.

The Howard Hughes Medical Institute, Sidney Kimmel Comprehensive Cancer Center, Johns Hopkins Medical Institutions, Baltimore, Maryland, USA.

Tumor endothelial marker 7 (TEM7) was recently identified as an mRNA transcript overexpressed in the blood vessels of human solid tumors. Here, we identify several new variants of TEM7, derived by alternative splicing, that are predicted to be intracellular (TEM7-I), secreted (TEM7-S), or on the cell surface membrane (TEM7-M) of tumor endothelium. Using new antibodies against the TEM7 protein, we confirmed the predicted expression of TEM7 on the cell surface and demonstrated that TEM7-M protein, like its mRNA, is overexpressed on the endothelium of various tumor types. We then used an affinity purification strategy to search for TEM7-binding proteins and identified cortactin as a protein capable of binding to the extracellular region of both TEM7 and its closest homologue, TEM7-related (TEM7R), which is also expressed in tumor endothelium. The binding domain of cortactin was mapped to a unique nine-amino acid region in its plexin-like domain. These studies establish the overexpression of TEM7 protein in tumor endothelium and provide new opportunities for the delivery of therapeutic and imaging agents to the vessels of solid tumors.

PMID: 15574754 [PubMed - indexed for MEDLINE]

Show:

[Write to the Help Desk](#)

[NCBI](#) | [NLM](#) | [NIH](#)

[Department of Health & Human Services](#)

[Privacy Statement](#) | [Freedom of Information Act](#) | [Disclaimer](#)